



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

At the vertex, $x'=0$, $y'=0$, and (2) becomes

$$x^2 + y^2 - px = 0 \dots\dots\dots (3).$$

If (x_1, y_1) be any point on the required locus, its polar with respect to (1) is

$$px - 2y_1y + px_1 = 0 \dots\dots\dots (4).$$

The condition that (4) touches (3) is

$$x_1^2 - y_1^2 - px_1 = 0 \dots\dots\dots (5),$$

an equilateral hyperbola.

II. Solution by J. SCHEFFER, A. M., Hagerstown, Maryland; CHAS. C. PURYEAR, Professor of Mathematics, Agricultural and Mechanical College, College Station, Texas; and G. B. M. ZERR, Texarkana, Ark.

Let $y^2 = 4ax$ be the equation to the parabola, (b, c) any point.

Then $cy = 2a(x + b) \dots\dots (1)$ is the polar of (b, c) . $x^2 + y^2 = ax \dots\dots (2)$ is the circle of curvature at the vertex. The value of y from (1) in (2) gives

$$c^2x^2 + 4a^2x^2 + 8a^2bx + 4a^2b^2 = ac^2x \dots\dots\dots (3).$$

From (3) we find the condition that (1) should be tangent to (2) to be

$$a^2c^4 = 8a^3bc^2 + 16a^2b^2c^2. \therefore c^2 = 8ab + 16b^2.$$

$$\therefore a^2 = (4b + a)^2 - c^2 \dots\dots\dots (4).$$

(4) represents an equilateral hyperbola.

CALCULUS.

Conducted by J. M. COLAW, Monterey, Va. All contributions to this department should be sent to him.

SOLUTIONS OF PROBLEMS.

55. Proposed by GEORGE LILLEY, Ph. D., LL. D., Principal of Park School, 394 Hall Street, Portland, Oregon.

A horse is tethered by a rope, a feet long, fastened to a post in a circular fence enclosing a circular piece of ground b feet in diameter. If the horse is tethered outside of the fence over how much ground can he feed? If he is inside the fence over how much ground can he feed? b is greater than a in each case.

$$\frac{d\theta}{d\rho} = \frac{4\rho}{b\sqrt{4\rho^2 - b^2}} - \frac{b}{2\rho^2 \sqrt{1 - \frac{b^2}{4\rho^2}}} = \frac{\sqrt{4\rho^2 - b^2}}{b\rho}.$$

$$\frac{dA}{d\rho} = \frac{dA}{d\theta} \frac{d\theta}{d\rho}, = \frac{\rho^2}{2} \times \frac{\sqrt{4\rho^2 - b^2}}{b\rho}, = \frac{\rho\sqrt{4\rho^2 - b^2}}{2b}.$$

Limits of ρ for *CFMA* are seen to be $\frac{1}{2}b$ and $\sqrt{a^2 + \frac{1}{4}b^2}$.

$$\therefore \text{Area } CFMA = \frac{1}{2b} \int_{\frac{1}{2}b}^{\sqrt{a^2 + \frac{1}{4}b^2}} \rho\sqrt{4\rho^2 - b^2} d\rho, = \frac{a^3}{3b}.$$

$$\text{Area } FHCAM = CFMA + C'CA - CFHC', = a^3/3b + ab/4 - ab/4 = a^3/3b.$$

$$\therefore \text{Area } F'BLAFHC' = 2a^3/3b + \pi a^2/2.$$

Internal area is composed of $2 \times \text{segment } PHC' + \text{sector } PDNC'$

$$\sin \frac{1}{2} \angle PCC' = (\frac{1}{2}a/\frac{1}{2}b) = a/b. \quad \angle PCC' = 2\sin^{-1}(a/b).$$

$$\angle PCN = 2\pi - 4\sin^{-1}(a/b). \quad \angle PC'N = \pi - 2\sin^{-1}(a/b).$$

$$\text{Sec. } PDNC' = \frac{a^2}{2} \left(\pi - 2\sin^{-1} \frac{a}{b} \right), \quad \text{sector } CPHC' = \frac{b^2}{8} \left(2\sin^{-1} \frac{a}{b} \right), = \frac{b^2}{4} \sin^{-1} \left(\frac{a}{b} \right)$$

$$\triangle PCC' = \frac{1}{2}a\sqrt{\frac{1}{4}b^2 - \frac{1}{4}a^2}, = \frac{1}{4}a\sqrt{b^2 - a^2}.$$

$$\text{Segment } PHC' = \frac{b^2}{4} \sin^{-1} \frac{a}{b} - \frac{1}{4}a\sqrt{b^2 - a^2}.$$

$$\therefore \text{Internal area} = \frac{a^2}{2} \left(\pi - 2\sin^{-1} \frac{a}{b} \right) + \frac{b^2}{2} \sin^{-1} \frac{a}{b} - \frac{1}{2}a\sqrt{b^2 - a^2},$$


$$= \frac{1}{2}\pi a^2 - \frac{a}{2}\sqrt{b^2 - a^2} + (\frac{1}{2}b^2 - a^2)\sin^{-1} \left(\frac{a}{b} \right).$$

III. Solution by G. B. M. YERR, A. M., Ph. D., Texarkana, Arkansas-Texas.

Let A be the point where the horse is tethered. $AF=a$, $AO=b/2$. Area $EADGKFE = 2 \text{ area } EAF + \text{area of semicircle } GKF$.

$$\therefore A = \int \rho^2 d\theta + \frac{1}{2}\pi a^2; \text{ but } \rho = \frac{1}{2}b\theta.$$

$$\therefore A = \frac{1}{2}b^2 \int_0^{2a/b} \theta^2 d\theta + \frac{1}{2}\pi a^2, = \frac{a^2}{6b}(4a + 3\pi b).$$



Let $x^2 + y^2 = \frac{1}{4}b^2$, be the equation to circle center O . $(x - \frac{1}{2}b)^2 + y^2 = a^2$, be the equation to circle center A .

$$\therefore OH = \frac{b^2 - 2a^2}{2b}, \quad \therefore BH = \frac{a}{b} \sqrt{b^2 - a^2}.$$

A' = area of segment BLC + area of segment BAC ,

$$\begin{aligned} &= \frac{b^2}{4} \left\{ \sin^{-1} \left(\frac{2a}{b^2} \sqrt{b^2 - a^2} \right) - \frac{2a(b^2 - 2a^2) \sqrt{b^2 - a^2}}{b^4} \right\} \\ &\quad + a^2 \left\{ \sin^{-1} \frac{\sqrt{b^2 - a^2}}{b} - \frac{a \sqrt{b^2 - a^2}}{b^2} \right\}, \\ &= \frac{1}{4} b^2 \cos^{-1} \left(\frac{b^2 - 2a^2}{b^2} \right) + a^2 \cos^{-1} \frac{a}{b} - \frac{a}{2} \sqrt{b^2 - a^2}. \end{aligned}$$


Also solved by J. SCHEFFER and A. H. HOLMES.

56. Proposed by B. F. BURLESON, Oneida Castle, New York.

Find (1) the length s of the closed curve of the cardioid; (2) its area A ; (3) if made to revolve about its axis $2a$, find the maximum longitudinal circumference C of the solid generated; (4) find the surface K of the same; (5) its volume V ; (6) the distance x_0 of the center of gravity of the solid from the origin O ; and (7) the distance g_0 of the center of gravity of the plane curve from the origin O .

I. Solution by J. SCHEFFER, A. M., Hagerstown, Maryland, and the PROPOSER.

Let $AB = a$ be the diameter of a circle. From A draw any chord AC . Make CP and $CP' = b$, then will the locus of P or P' be the Limaçon. If $AP = r$, $\angle PAB = \theta$, we find at once the polar equation of the Limaçon to be $r = a \cos \theta + b$. If $b > a$, the curve consists of but one loop; if $b < a$, it has two loops, and if $b = a$ the curve becomes the Cardioid, the polar equation of which is $r = a(1 + \cos \theta)$. It can easily be shown that the cardioid is an epicycloid, the generating circle of which is equal to the fixed one; also, drawing through the center O of the circle a line parallel to AP cutting the circumference of the circle at D , and drawing through P a line parallel to CD , this line is a tangent to the cardioid at P . The different problems proposed are best solved by means of the polar equation of the curve.



$$(1). \quad \text{The length } s = 2 \int_0^\pi d\theta \sqrt{r^2 + \left(\frac{dr}{d\theta} \right)^2} = 2a \int_0^\pi \cos \frac{1}{2} \theta d\theta = 8a.$$

$$(2). \quad \text{The area } A = \int_0^\pi r^2 d\theta = a^2 \int_0^\pi (1 + \cos \theta)^2 d\theta = \frac{3}{2} \pi a^2.$$